

# WARNING

Keep out of the reach of children. Read the label before use.

# FERACOL®

Contains cholecalciferol 8g/kg in the form of a bait.

## FOR THE CONTROL OF POSSUMS AND RODENTS

**WARNING:** may be harmful if swallowed. Suspected of damaging fertility or the unborn child. Harmful to aquatic Life.

**PRECAUTIONS:** keep out of reach of children and away from domestic animals and foodstuffs. Avoid contact with skin. Wear gloves. Wash hands after use and before eating, drinking or smoking. If exposed or concerned: Get medical advice. Exclude dogs from baiting area as they are very susceptible to cholecalciferol.

**DIRECTIONS FOR USE:** Feracol paste has flavour and lure incorporated during manufacture, no further additives are required. Use only in bait stations or bait bags. Maintain an uninterrupted supply of bait until signs of activity ceases. Only a single feed of Feracol paste is required, the onset of symptoms prevents further feeding of toxic baits. It is critical that sufficient bait is applied to ensure all animals visiting the bait will receive a toxic dose during one feeding session to help prevent sub-lethal poisoning and the development of aversion. A pre-feed with non-toxic Ferafeed paste is strongly recommended to maximise toxic bait uptake. After pre-feeding replace non-toxic Ferafeed paste with Feracol paste. A judgement of pest abundance is required to gauge the amount of bait to apply.

**Possum control:** a lethal dose for a possum is 18g (approximately a dessert spoon). Add at least this amount to each bait bag or bait station.

**Rodent control:** a lethal dose for a rat is about 3g but depends on bodyweight. For best results use 40g Feracol paste in bait stations placed at 50m intervals in a grid pattern covering the entire area to be treated. Check every 4 days and replace baits where required. If possible avoid toxic treatments during breeding peaks.

**SYMPTOMS OF POISONING** no symptoms may occur for several hours or days after poisoning has occurred. Symptoms may include loss of appetite, thirst, diarrhoea, possible trembling and lethargy.

**FIRST AID:** in the event of poisoning or suspected poisoning do not induce vomiting. Call a doctor or take to nearest hospital without delay. Show this label or safety data sheet. For further advice contact National Poison Centre 0800 POISON (0800 764 766). For animals, consult a veterinarian immediately.

**DOMESTIC ANIMAL/LIVESTOCK:** when used in the vicinity of domestic animals ensure they cannot access the bait and the bait is in a secure bait station. Feracol is very hazardous to cats and dogs. It is important to remove and prevent access to bait contents. Owners of pets and animals in the immediate vicinity should be notified of this risk if these pets or animals may be impacted. Stock must be kept out of treated areas until uneaten bait is removed or destroyed.

**LEGAL OBLIGATIONS:** by law this product must only be used in bait stations or bags. When the product is used in a place to which

members of the public ordinarily have access, signs must be erected at every normal point of entry to warn the public that this product has been applied in the area. Signs must remain in place for not less than 4 months from the last date on which baits were laid if biodegradable bait bags were used, or 2 to 3 months after baits have been retrieved from the place where it has been laid if bait stations are used. The product must only be used as specified on the label. Feracol is not to be used in households or industrial buildings.

**SHELF LIFE:** any product held after the expiry date shown on the container should be disposed of according to the label directions.

**STORAGE:** Store in a cool place where temperature does not exceed 25°C. Store in a tightly closed container. Store locked up.

**SPILLAGE:** in the event of a spill, isolate the spill area and take all practical steps to manage any harmful effects. Scoop up baits into secure labelled containers. Undamaged bait may be reused. Prevent split bait entering streams or waterways. Dispose of spoiled bait according to the label directions.

**DISPOSAL:** The potency of the bait will reduce with age, UV light, heat and microbial action. Dispose of unwanted or expired product by burying under at least 60cm of damp biologically active soil or with other organic matter in an active appropriately managed landfill. Ensure bait is covered immediately to prevent scavenging. Unwanted or expired product can be burnt in a suitably constructed and located incinerator and any residues buried as above. Do not use the empty container for anything else. Dispose of empty containers by burying or burning as described.

**CONDITIONS OF SALE:** Aotea IP Holdings Ltd and Connovation Ltd will not accept any responsibility for any consequential loss, injury or damage arising in connection with supply or use of these goods other than the goods be of merchantable quality at the time of sale. To the extent allowed by such statutes the liability of Aotea IP Holdings Ltd and Connovation Ltd is limited to the replacement or refund of the goods at the price paid.

**HSNO Approval Code:** HSR001598 HSNO  
**HSNO Classifications:** 6.1E (O), 6.8B, 9.1D  
Registered pursuant to the ACVM Act 1997 No. V5263  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

Batch No: .....

Label No: .....

Date of Expiry: .....

Bagged Paste: ☐ 500

Manufactured & Distributed by:

FeraCol® is the registered trademark of Aotea IP Holdings Ltd



36B Sir William Ave, East Tamaki, Manukau, Auckland  
T: +64 9 273 4333 | M: +64 9 273 4888 | [www.connovation.co.nz](http://www.connovation.co.nz)

